

# Booster Neutrino Experiment


Alexis Aguilar–Arevalo  
Columbia University  
MiniBooNE Collaboration

Neutrino Planning Meeting  
Santa Fe, New Mexico  
October 29, 2005

# The MiniBooNE experiment and LSND

MiniBooNE was designed to confirm or refute the LSND oscillations signal

## The LSND oscillations signal



Observed excess:


**87.9 ± 22.4 ± 6.0**

$\nu_e$  events

Oscillation

Probability:

**0.264 ± 0.067 ± 0.045 %**



LSND + Karmen2  
combined analysis  
allowed region

**BooNE:** a 2<sup>nd</sup> generation experiment that will come to life if MiniBooNE confirms LSND. MiniBooNE design allows to build on to BooNE.


# Mini-BooNE Collaboration




**MiniBooNE consists of about 70  
scientists from 14 institutions.**

Y. Liu, D. Perevalov, I. Stancu *Alabama*  
S. Koutsoliotas *Bucknell*  
E. Hawker, R.A. Johnson, J.L. Raaf *Cincinnati*  
T. Hart, R.H. Nelson, E.D. Zimmerman *Colorado*  
A. Aguilar-Arevalo, L. Bugel, L. Coney, J.M. Conrad,  
J. Link, J. Monroe, K. Mahn, D. Schmitz,  
M.H. Shaevitz, G.P. Zeller, Z. Djurcic *Columbia*  
D. Smith *Embry Riddle*  
L. Bartoszek, C. Bhat, S. J. Brice, B.C. Brown,  
D.A. Finley, R. Ford, F.G. Garcia,  
P. Kasper, T. Kobilarcik, I. Kourbanis,  
A. Malensek, W. Marsh, P. Martin, F. Mills,  
C. Moore, P. Nienaber, E. Prebys,  
A.D. Russell, P. Spentzouris, R. Stefanski,  
T. Williams *Fermilab*  
D. C. Cox, A. Green, H.-O. Meyer, R. Tayloe, T. Kattori  
*Indiana*  
G.T. Garvey, C. Green, W.C. Louis, G. McGregor,  
S. McKenney, G.B. Mills, V. Sandberg,  
B. Sapp, R. Schirato, R. Van de Water,  
D.H. White *Los Alamos*  
R. Imlay, W. Metcalf, M. Sung, M.O. Wascko  
*Louisiana State*  
J. Cao, Y. Liu, B.P. Roe, H. Yang *Michigan*  
A.O. Bazarko, P.D. Meyers, R.B. Patterson,  
F.C. Shoemaker, H.A. Tanaka *Princeton*  
M. Sorel, *U. Valencia*  
B.T. Flemming, A. Curionni, *Yale*

# MiniBooNE Neutrino Beam




8 GeV Proton Beam Transport



50m Decay Pipe


Variable decay pipe length  
(2 absorbers @ 50m and 25m)



One magnetic horn, with Be target





Magnetic Horn




Detector

# The MiniBooNE Detector




- 12 meter diameter sphere
- Filled with 950,000 liters (800 tons) of very pure mineral oil
- Light tight inner region with 1280 photomultiplier tubes
- Outer veto region with 241 PMTs.
- **Oscillation Search Method:**

Look for  $\nu_e$  events  
in a pure  $\nu_\mu$  beam




# Particle Identification

- Separation of  $\nu_\mu$  from  $\nu_e$  events
  - Exiting  $\nu_\mu$  events fire the veto
  - Stopping  $\nu_\mu$  events have a Michel electron after a few  $\mu\text{sec}$
  - Also, scintillation light with longer time constant  $\Rightarrow$  enhanced for slow pions and protons
  - Čerenkov rings from outgoing particles
    - Shows up as a ring of hits in the phototubes mounted inside the MiniBooNE sphere
    - Pattern of phototube hits indicates the particle type



# Intrinsic $\nu_e$ in the beam

Small intrinsic  $\nu_e$  rate  $\Rightarrow$  Event Ratio  $\nu_e/\nu_\mu = 6 \times 10^{-3}$



$\nu_e$  from  $\mu$ -decay



- Directly tied to the observed half-million  $\nu_\mu$  interactions



Kaon rates measured in low energy proton production experiments


- HARP experiment (CERN)
- E910 (Brookhaven)
- MiniBooNE “High Energy Box” data

- “Little Muon Counter” measures rate of kaons in-situ





# Estimates for the $\nu_\mu \rightarrow \nu_e$ Appearance Search

- Look for appearance of  $\nu_e$  events above background expectation
  - Use data measurements both internal and external to constrain background rates





- Fit to  $E_\nu$  distribution used to separate background from signal.



# MiniBooNE Oscillation Sensitivity

- Oscillation sensitivity and measurement capability
  - Data sample corresponding to  $1 \times 10^{21}$  pot
  - Systematic errors on the backgrounds average  $\sim 5\%$




## If MiniBooNE sees a signal...

- \* If MiniBooNE sees no indications of oscillations with  $\nu_\mu$ 
  - $\Rightarrow$  **Need to run with  $\bar{\nu}_\mu$  since LSND signal was  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$**   
(Morgan O. Wascko, PANIC'05)
- If MiniBooNE sees an oscillation signal
  - $\Rightarrow$  **There are 3  $\Delta m^2$  scales involved in oscillations**

How can there be three distinct  $\Delta m^2$ 's?

- Additional “sterile” neutrinos involved in oscillations
- One of the experimental measurements is not neutrino oscillations
  - [e.g. Neutrino decay (Palomares–Ruiz *et.al.*, hep-ph/0505216) ]
- CP violation and sterile  $\nu$ 's (allows different mixing for  $\nu$ 's and anti- $\nu$ 's)
- Even stranger things (CPT violation,...)

# MiniBooNE possibilities -- oscillations



# Experimental Program with Sterile Neutrinos

If sterile neutrinos is the answer, then many mixing angles, CP phases, and  $\Delta m^2$  could be included

- Measure number of extra masses  $\Delta m_{14}^2, \Delta m_{15}^2 \dots$
- Measure mixings  
Could be many small angles

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \\ \nu_s \\ \nu_{s'} \\ \vdots \end{pmatrix} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} & U_{e4} & U_{e5} & \dots \\ U_{\mu1} & U_{\mu2} & U_{\mu3} & U_{\mu4} & U_{\mu5} \\ U_{\tau1} & U_{\tau2} & U_{\tau3} & U_{\tau4} & U_{\tau5} \\ U_{s1} & U_{s2} & U_{s3} & U_{s4} & U_{s5} \\ U_{s'1} & U_{s'2} & U_{s'3} & U_{s'4} & U_{s'5} \\ \dots & & & & & \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \\ \nu_4 \\ \nu_5 \\ \vdots \end{pmatrix}$$



Map out mixings associated with  $\nu_\mu \rightarrow \nu_e$

Map out mixings associated with  $\nu_\mu \rightarrow \nu_\tau$

- Oscillations to sterile neutrinos could affect long-baseline measurements and strategy
- Compare  $\nu_\mu$  and  $\bar{\nu}_\mu$  oscillations  $\Rightarrow$  CP and CPT violations

# If MiniBooNE sees $\nu_\mu \rightarrow \nu_e$ then: BooNE: Two Detector Exp.

- Far detector at 1.025 km for high  $\Delta m^2$  signal or 2.025km for low  $\Delta m^2$  signal.
- Cost of one additional detector ranges from 5 to 10 Million Dollars (conceivable to think about more than one)




- Precision measurement of oscillation parameters
  - $\sin^2 2\theta$  and  $\Delta m^2$
  - Map out the  $n \times n$  mixing matrix
- Determine how many high mass  $\Delta m^2$  's
  - 3+1, 3+2, 3+3 ...
- Show the L/E oscillation dependence
  - Oscillations or  $\nu$  decay or ???

## Two location BooNE experiment:

1 near detector at current MiniBooNE location and 1 far detector at 1025m, (2025m) if observe a signal with oscillation parameters:  $[\sin^2 2\theta, \Delta m^2] = [0.4 \text{ eV}^2, 0.017]$  **Low  $\Delta m^2$**   
 $[\sin^2 2\theta, \Delta m^2] = [1.0 \text{ eV}^2, 0.004]$  **High  $\Delta m^2$**


# BooNE: 1<sub>near</sub>+ 4<sub>far</sub> Detector Exp., and other possibilities

Much higher precision can be achieved by adding more detectors (e.g. 4- det)



Two location BooNE experiment:  
1 near detector at current MiniBooNE location and 4 far detectors at second location. Same correspondence to signals as in previous slide.

Also explore disappearance in high  $\Delta m^2$  to probe oscillations into sterile  $\nu$ 's.





A near detector at  $\sim 100\text{m}$  (Finesse proposal) for disappearance and background determination could be useful (See R.Tayloe's talk)

If MiniBooNE sees  $\nu_\mu \rightarrow \nu_e$  (or not) then:  
 run MiniBooNE with anti-neutrinos for  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$

- Are  $\nu_\mu$  and  $\bar{\nu}_\mu$  the same?
  - Mixing angles,  $\Delta m^2$  values
- Explore CP (or CPT) violation by comparing  $\nu_\mu$  and  $\bar{\nu}_\mu$  results
- Running with anti-neutrinos takes longer to obtain similar sensitivity

[M.O. Wascko, PANIC'05]



MiniBooNE appearance sensitivity region for anti-neutrino oscillations in case of no oscillation in neutrinos

If we see a signal in anti-neutrinos:  
 run BooNE with anti-neutrinos for  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$

## A possible time scale for BooNE (2 detector system)

- 2006 appearance results and anti- $\nu$  running
- 3 years after signal is observed start detector construction.
- ~2 years later begin data taking with 2-detector system

### Things to note:

-----

- \* Detector identical in design to MiniBooNE (systematics motivated)
- \* Would use:
  - New Oil
  - New electronics (Los Alamos developing design)
- \* Construction timescales and costs well understood
  - Estimated 5-7 MD for one additional Detector
- \* WBS already in place

Ready to go in the event of a positive MiniBooNE result!

## Conclusions




- A positive MiniBooNE result will bring us to a new and exciting era in neutrino physics.
- Additional detector(s) can boost the measurement capabilities of the booster neutrino beam at Fermilab and make it unique in its kind.
- Precision measurements will allow us to test CP conservation in neutrino sector with the existing beamline and its capability of change in polarity.
- The direction of the field will be determined by what we discover in these experiments

# Backup Slides




## $\nu_\mu \rightarrow \nu_\tau$ Appearance at High $\Delta m^2$

- Appearance of  $\nu_\tau$  helps to sort out the mixings through the sterile components
- Need moderately high neutrino energy to get above the 3.5 GeV  $\tau$  threshold (~6-10 GeV)
- Example: NuMI Med energy beam 8 GeV with detector at L=2km (116m deep)

Emulsion Detector or Liquid Argon



# Explain LSND with Sterile Neutrinos



- Constraints from the atmospheric and solar data indicate:

⇒ Sterile neutrino is mainly associated with the LSND  $\Delta m^2$




Then these are the main mixing matrix elements

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \\ \nu_s \\ \nu_{s'} \\ \vdots \end{pmatrix} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} & U_{e4} & U_{e5} & \dots \\ U_{\mu1} & U_{\mu2} & U_{\mu3} & U_{\mu4} & U_{\mu5} & \dots \\ U_{\tau1} & U_{\tau2} & U_{\tau3} & U_{\tau4} & U_{\tau5} & \dots \\ U_{s1} & U_{s2} & U_{s3} & U_{s4} & U_{s5} & \dots \\ U_{s'1} & U_{s'2} & U_{s'3} & U_{s'4} & U_{s'5} & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \\ \nu_4 \\ \nu_5 \\ \vdots \end{pmatrix}$$

Check via  
 $\nu_e$  appearance and  
 $\nu_\mu$  disappearance

# CPT Violation in the neutrino sector ( $m_\nu \neq m_{\bar{\nu}}$ )

*Accommodates all three signals with three standard model neutrinos*



hep-ph/0210411  
Barenboim, Lykken

- A new non-local field theory
  - not Lorentz violating
  - introducing a whole new "Dirac Eq."
- Fit all present data, including Super K and KamLand

*Check by comparing neutrino and anti-neutrino modes*